

## 引用文獻

### 一、中文部份

江昶勳 (2007)。探討不同類型之馬拉松對人體壓力荷爾蒙及生理負荷相關指標的影響。未出版碩士論文，國立陽明大學，台北市。

吳輝秋 (2006)。認識輔酶 Q<sub>10</sub>。高雄縣醫師會誌，22，11-16。

李小龍 (2004)。輔酶 Q<sub>10</sub> 對耐力型運動員最大運動能力及乳酸閾值 (OBLA) 之影響。未出版碩士論文，中國文化大學，台北市。

李正華 (1999)。心肌損傷生化學標幟。當代醫學，26 (12)，46-50。

李春璟 (2004)。液相層析法對人體血漿中 Coenzyme Q<sub>10</sub> 之分析及應用。未出版碩士論文，高雄醫學大學，高雄市。

李建明、黃欽永、黃谷臣、湯馥君 (2002)。不同年齡層參與馬拉松的血液生化值之比較。大專體育學刊，12，157-164。

李聰明 (1998)。Troponin T/I 及其臨床意義。基層醫學，12 (13)，234-237

吳慧君 (2004)。24 小時超級馬拉松跑對選手血、尿液生化值之影響。台北市：文化大學出版部

吳慧君 (1999)。運動能力的生理學評定。台北市：師大書院。

林文發 (2000)。運動訓練生化分析。台北市：文化大學出版部。

林貴福 (1997)。無氧能量代謝酵素乳酸去氫酶研究的分析。台灣師大體育研究，3，121-136。

邱榮基、蔚順華 (2005)。馬拉松長跑選手的運動猝死。大專體育，76，210-214。

陳慧汝 (1996)。Melatonin 及 Magnolol 對大鼠心肌缺血一再循環損傷之保

護作用。未出版碩士論文，國防醫學院，台北市。

陳忠慶、陳信良（2005）。離心運動對血液肌肉蛋白質評估指標的反應。運動生理暨體能學報，2，1-17。

溫雅萍（2002）。血漿裂解肌肉蛋白-Troponin I 影響因子之研究。未出版碩士論文，國立中興大學，台中市。

賴紫蘭、吳慧君（2005）。不同方式輔酶 Q<sub>10</sub> 補充對於籃球運動員有氧能力及運動後恢復其生理生化之影響。大專體育學刊，1（7），269-278。



## 二、英文部分

- Ammann, P., Pfisterer, M., Fehr, T., & Rickli, H. (2004). Raised cardiac troponins. *British Medical Journal*, 328, 1028-1029.
- Battino, M., Fato, R., Parenti-Castelli, G., & Lenaz, G. (1990). Coenzyme Q can control the efficiency of oxidative phosphorylation. *International Journal Tissue Reactive*, 12(3), 137-144.
- Beyer, R. E. (1992). An analysis of the role of coenzyme Q in free radical generation and as an antioxidant. *Biochemistry Cell and Biology*, 70(6), 390-403.
- Belardinelli, R., Muçaj, A., Lacalaprice, F., Solenghi, M., Principi, F., Tiano, L., et al. (2005). Coenzyme Q10 improves contractility of dysfunctional myocardium in chronic heart failure. *Biofactors*, 25(1-4), 137-145.
- Bodor, G. S., Porterfield, D., Voss, E. M., Smith, S., & Apple, F. S. (1995). Cardiac troponin-I is not expressed in fetal and healthy or diseased adult human skeletal muscle tissue. *Clinical Chemistry*, 41(12 Pt 1), 1710-1715.
- Chosa, E., Sekimoto, T., Sonoda, N., Yamamoto, K., Matsuda, H., Takahama, K., et al. (2003). Evaluation of human beta-enolase as a serum marker for exercise-induced muscle damage. *Clinical Journal of Sport Medicine*, 13(4), 209-212.
- Chen, T. C., & Hsieh, S. S. (2000). The effects of repeated maximal voluntary isokinetic eccentric exercise on recovery from muscle damage. *Research Quarterly for Exercise and Sport*, 71, 260-266.
- Chew, G. T., & Watts, G. F. (2004). Coenzyme Q10 and diabetic endotheliopathy: oxidative stress and the 'recoupling hypothesis'. *Monthly Journal of the Association of Physicians*, 97(8), 537-48.
- Chapelle, J. P. (1999). Cardiac troponin I and troponin T: recent plays in the

filed of myocardial makers. *Clinical Chemistry and Laboratory Medicine*, 37(1), 11-20.

Ernster, L., & Dallne G. (1995). Biochemical, physiological and medical aspects of ubiquinone function. *Biochimica et Biophysica Acta*, 1271(1), 195-204.

Fallon, K, E., Sivyer, G., Sivyer, K., & Dare, A. (1999). The biochemistry of runners in a 1600 km ultramarathon. *British Journal of Sports Medicine*, 33(4), 264-269.

Fornes, P., & Lecomte, D. (2003). Pathology of sudden death during recreational sports activity: an autopsy study of 31 cases. *The American Journal of Forensic Medicine and Pathology*, 24(1), 9-16.

Futterman , L. G., & Myerburg, R. (1998). Sudden death in athletes: an update. *Sports Medicine*, 26(5), 335-350.

Greenberg, S., & Frishman, W. H. (1990). Co-enzyme Q10: a new drug for cardiovascular disease. *Journal of Clinical Pharmacology*, 30(7), 596-608.

Hoogsteen, J., Hoogeveen, A., Schaffers, H., Wijn, P.F., van Hemel, N.M., & van der Wall, E. E. (2004). Myocardial adaptation in different endurance sports: an echocardiographic study. *The International Journal of Cardiovascular Imaging*, 20(1), 19-26.

Ikematsu, H., Nakamura, K., Harashima, S., Fujii, K., & Fukutomi, N. (2006). Safety assessment of coenzyme Q10 (Kaneka Q10) in healthy subjects: a double-blind, randomized, placebo-controlled trial. *Regulatory Toxicology and Pharmacology*, 44(3), 212-218.

Ishikawa, Y., Saffitz, J. E., Mealman, T. L., Grace, A. M., & Roberts, R. (1997). Reversible myocardial ischemic injury is not associated with increased creatine kinase activity in plasma. *Advances in Clinical Chemistry*, 43(3), 467-75.

- Ito, H., Nakajima, T., Takikawa, R., Hamada, E., Iguchi, M., Sugimoto, T., et al. (1991). Coenzyme Q10 attenuates cyanide-activation of the ATP-sensitive K<sup>+</sup> channel current in single cardiac myocytes of the guinea-pig. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 344, 133-136.
- Katus, H. A., Looser, S., Hallermayer, K., Remppis, A., Scheffold, T., Borgya, A., et al. (1992). Development and vitro characterization of new immunoassay of cardiac troponin T. *Advances in Clinical Chemistry*, 38, 386-393.
- Katus, H. A., Remppis, A., Scheffold, T., Diederich, K. W., & Kuebler, W. (1991). Intracellular compartmentation of cardiac troponin T and its release kinetics in patients with reperfused and nonreperfused myocardial infarction. *The American Journal of Cardiology*, 67, 1360-1367.
- Kiely, P. D., Bruckner, F. E., Nisbet, J. A., & Dagher, A. (2000). Serum skeletal troponin I in inflammatory muscle disease: relation to creatine kinase, CKMB and cardiac troponin I. *Annals of the Rheumatic Diseases*, 59(9), 750-751.
- Kishimoto, C., Tomioka, N., Nakayama, Y., & Miyamoto, M. (2003). Anti-oxidant effects of coenzyme Q10 on experimental viral myocarditis in mice. *Journal of cardiovascular pharmacology*, 42(5), 588-592.
- Kobayashi, Y., Takeuchi, T., Hosoi, T., Yoshizaki, H., & Loeppky, J. A. (2005). Effect of a marathon run on serum lipoproteins, creatine kinase, and lactate dehydrogenase in recreational runners. *Research Quarterly for Exercise and Sport*, 76(4), 450-455.
- Kokado, H., Shimizu, M., Yoshio, H., Ino, H., Okeie, K., Emoto, Y., et al. (2000). Clinical features of hypertrophic cardiomyopathy caused by a Lys 183 deletion mutation in the cardiac troponin I gene. *Circulation*, 102, 663-669.

- König, D., Schumacher, Y. O., Heinrich, L., Schmid, A., Berg, A., & Dickhuth, H. H. (2003). Myocardial stress after competitive exercise in professional road cyclists. *Medicine and Science in Sports and Exercise*, 35(10), 1679-1683.
- Kratz, A., Lewandrowski, K. B., Siegel, A. J., Chun, K. Y., Flood, J. G., Van Cott, E. M., et al. (2002). Effect of marathon running on hematologic and biochemical laboratory parameters, including cardiac markers. *American Journal of Clinical Pathology*, 118(6), 856-863.
- La Gerche, A., Boyle, A., Wilson, A. M., & Prior, D. L. (2004). No evidence of sustained myocardial injury following an Ironman distance triathlon. *International Journal of Sports Medicine*, 25(1), 45-49.
- Lenaz, G. (2001). A critical appraisal of the mitochondrial coenzyme Q pool. *FEBS Letters*, 509(2), 151-155.
- Lockwood, K., Moesgaard, S., Hanioka, T., & Folkers, K. (1994). Apparent partial remission of breast cancer in 'high risk' patients supplemented with nutritional antioxidants, essential fatty acids and coenzyme Q10. *Molecular Aspects of Medicine*, 15, 231-240.
- Lampertico, M., & Comis, S. (1993). Italian multicenter study on the efficacy and safety of coenzyme Q10 as adjuvant therapy in heart failure. *The Clinical Investigator*, 71(8 Suppl), 129-133.
- Lenaz, G., Fato, R., Castelluccio, C., Genova, M. L., Bovina, C., Estornell, E., et al. (1993). The function of coenzyme Q in mitochondria. *The Clinical Investigator*, 71(8 Suppl), 66-70.
- Leavis, P. C., & Gergely, J. (1984). Thin filament proteins and thin filament-linked regulations of vertebrate muscle contraction. *CRC Critical Reviews in Biochemistry*, 16, 235-305.
- Margaritis, I., Tessier, F., Verdera, F., Bermon, S., & Marconnet, P. (1999).

- Muscle enzyme release does not predict muscle function impairment after triathlon. *The Journal of Sports Medicine and Physical Fitness*, 39(2), 133-139.
- Maron, B. J., Shirani, J., Poliac, L. C., Mathenge, R., Roberts, W. C., & Mueller, F. O. (1996). Sudden death in young competitive athletes. Clinical, demographic, and pathological profiles. *The Journal of the American Medical Association*, 276, 199-204.
- Mena, P., Maynar, M., & Campillo, J. E. (1996). Changes in plasma enzyme activities in professional racing cyclists. *British Journal of Sports Medicine*, 30(2), 122-124.
- Missov, E., Calzolari, C., & Pau, B. (1997). Circulating cardiac troponin I in severe congestive heart failure. *Circulation*, 96, 2953-2958.
- Mortensen, S. A. (1993). Perspectives on therapy of cardiovascular diseases with coenzyme Q10 (ubiquinone). *The Clinical Investigator*, 71(8 Suppl), 116-123.
- Mortensen, S. A. (2003). Overview on coenzyme Q10 as adjunctive therapy in chronic heart failure. Rationale, design and end-points of "Q-symbio" -a multinational trial. *Biofactors*, 18(1-4), 79-89.
- Nohl, H., Kozlov, A. V., Staniek, K., & Gille, L. (2001). The multiple functions of coenzyme Q. *Bioorganic Chemistry*, 29(1), 1-13.
- Nosaka, K., & Clarkson, M. P. (1995). Muscle damage following repeated bouts of high force eccentric exercise. *Medicine and Science in Sports and Exercise*, 27, 1263-1269.
- Pobezhimova, T. P., & Voinikov, V. K. (2000). Biochemical and physiological aspects of ubiquinone function. *Membrane & Cell Biology*, 13(5), 595-602.

- Rosenfeldt, F., Hilton, D., Pepe, S., & Krum, H. (2003). Systematic review of effect of coenzyme Q10 in physical exercise, hypertension and heart failure. *Biofactors*, 18(1-4), 91-100.
- Rosenfeldt, F., Marasco, S., Lyon, W., Wowk, M., Sheeran, F., Bailey, M., et al. (2005). Coenzyme Q10 therapy before cardiac surgery improves mitochondrial function and in vitro contractility of myocardial tissue. *The Journal of Thoracic and Cardiovascular Surgery*, 129(1), 25-32.
- Robert, W. O. (1998). Medical supervision of the athlete-event coverage. *Manual of Sports Medicine* (pp. 54-65). MR Safran, Lippincott-Raven Publishers: Philadelphia.
- Salles, J. E., Moises, V. A., Almeida, D. R., Chacra, A. R., & Moises, R. S. (2005). Myocardial dysfunction in mitochondrial diabetes treated with Coenzyme Q10. *Diabetes Research and Clinical Practice*, 72(1), 100-103.
- Shave, R. E., Dawson, E., Whyte, G., George, K., Gaze, D., & Collinson, P. (2004). Effect of prolonged exercise in a hypoxic environment on cardiac function and cardiac troponin T. *British Journal of Sports Medicine*, 38(1), 86-88.
- Shults, C.W. (2003). Coenzyme Q10 in neurodegenerative diseases. *Current Medicinal Chemistry*, 10(19), 1917-1921.
- Smith, J. E., Garbutt, G., Lopes, P., & Pedoe, D. T. (2004). Effects of prolonged strenuous exercise (marathon running) on biochemical and haematological markers used in the investigation of patients in the emergency department. *British Journal of Sports Medicine*, 38(3), 292-294.
- Shave, R. E., Dawson, E., Whyte, P. G., George, K., Ball, D., Gaze, C. D., et al.



- (2003). Cardiac troponin T in female athletes during a two-day mountain marathon. *Scottish Medical Journal*, 48(2), 41-42.
- Singh, R. B., Niaz, M. A., Rastogi, V., & Rastogi, S. S. (1998). Coenzyme Q in cardiovascular disease. *The Journal of the Association of Physicians of India*, 46(3), 299-306.
- Tran, M. T., Mitchell, T. M. Kennedy, D. T., & Giles, J. T. (2001). Role of coenzyme Q10 in chronic heart failure, angina, and hypertension. *Pharmacotherapy*, 21, 797-806.
- Triolo, L., Lippa, S., Oradei, A., De Sole, P., & Mori, R. (1994). Serum coenzyme Q10 in uremic patients on chronic hemodialysis. *Nephron*, 66(2), 153-156.
- Vidotto, C., Tschan, H., Atamaniuk, J., Pokan, R., Bachl, N., & Muller, M. M. (2005). Responses of N-terminal pro-brain natriuretic peptide (NT-proBNP) and cardiac troponin I (cTnI) to competitive endurance exercise in recreational athletes. *International Journal Sports Medicine*, 26(8), 645-650.
- Vuori, I. M. (1995). Sudden death and exercise: effects of age type of activity. *Scandinavian Journal of Medicine and Science in Sports*, 5(5), 267-268.
- Wu, C. J., Liang, H. L., Chiou, K. R., Mar, G. Y., Tseng, C. J., Lin, et al. (2001). Significance of Cardiac Troponin I and Creatine Kinase Release after Coronary Intervention. *Zhonghua Yi Xue Za Zhi (Taipei)*, 64, 343-350.
- Zhou, S., Zhang, Y., Davie, A., Marshall-Gradisnik, S., Hu, H., Wang, J., et al. (2005). Muscle and plasma coenzyme Q(10) concentration, aerobic power and exercise economy of healthy men in response to four weeks of supplementation. *Journal Sports Medicine Physical Fitness*, 45(3), 337-346.

Ziegler, S., Niessner, A., Slany, J., Muller, M. M., Heinschink, A., Billensteiner, E., et al. (2003). Muscle cell proteins are selectively released into the blood stream by marathon running. *Acta Medica Austriaca*, 30(2), 55-58.

Zot, A. S., & Pptter, T. D. (1987). Structural aspects of troponin-tropomyosin regulation of skeletal muscle contraction. *Annual Review Biophysics and Biophysical Chemistry*, 16, 535-559.

林正常等（譯）（2003）。*運動生理學：體適能與運動表現的理論與應用*。台北市：藝軒圖書出版社。（Scott, K., & Edward, E., 2001）

